

*Citation for published version:*

Feil, EJ 2012, 'The emergence and spread of dysentery', *Nature Genetics*, vol. 44, no. 9, pp. 964-965.  
<https://doi.org/10.1038/ng.2389>

*DOI:*

[10.1038/ng.2389](https://doi.org/10.1038/ng.2389)

*Publication date:*

2012

*Document Version*

Peer reviewed version

[Link to publication](#)

## University of Bath

### Alternative formats

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## The emergence and spread of dysentery

Edward J. Feil

*Professor Edward J. Feil is at the Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY UK. Email: e.feil@bath.ac.uk*

***Shigella sonnei* is an important cause of bacterial dysentery in the developed world, and has also recently emerged in transitional countries. Phylogenetic analysis based on whole-genome sequencing of a global sample has detailed the recent evolutionary history of this pathogen, and shed light on the genetic changes associated with this epidemiological shift.**

*Shigella* are human associated pathogenic *E. coli*, transmitted by the fecal-oral route, and are responsible for at least 120 million cases of dysentery each year, with approximately 1 million fatalities<sup>1</sup>. On page xx of this issue, Nicholas Thomson and colleagues<sup>2</sup> describe a detailed evolutionary analysis of a diverse collection of *Shigella sonnei* isolates. The work provides an exemplar of how next-generation sequencing has transformed our ability to reconstruct historical and contemporary patterns of transmission and genome divergence within bacterial pathogens<sup>3</sup>

Four *Shigella* serogroups have been described, each of which independently acquired a pINV plasmid conferring the ability to invade the gut mucosa<sup>4</sup>. These serogroups exhibit distinct epidemiological properties; for example, *S. flexneri* is waterborne and prevalent in the developing world, whereas *S. sonnei*, which is less diverse than other serogroups, more typically causes disease in the developed world via person-to-person spread or contaminated food<sup>5</sup>. Improved sanitation over last few decades has led to a decrease in *S. flexneri* cases in the developing world, but has coincided with the emergence of *S. sonnei*.

### Sequencing and phylogenetic analysis

Holt et al. assembled 132 *S. sonnei* strains isolated between 1943 and 2008 from four continents. This diverse sample allowed the inference of long-range transmission events and local expansion, while the inclusion of older isolates enabled reliable temporal calibration of the tree. The whole genomes of pools of 11 or 12 isolates were sequenced on the Illumina Genome Analyzer GAI, and the authors reconstructed robust maximum likelihood trees using RAxML with *E. coli* and *Shigella* outgroups. Given the tree topology and isolation dates it is possible to test for temporal signal in the data, and to date the major nodes in the tree. Holt et al. estimated a mean mutation rate of  $6 \times 10^{-7}$  site<sup>-1</sup> year<sup>-1</sup>, approximately half the rate of *S. aureus*<sup>6</sup>, but over twice as fast as the highly monomorphic *Yersinia pestis*<sup>7</sup>. This rate variation between taxa is most readily explained by longer generation times in more slowly evolving species.

Holt et al. estimate the most recent common ancestor of *S. sonnei* to have emerged under 500 years ago, far more recently than previously thought<sup>8</sup>. Four robust lineages are resolved from the data, one of which (lineage IV) is represented by a single isolate. These major divisions are consistent with *S. sonnei* biotypes, CRISPR

types, and are likely to be resolved by IST, MLVA<sup>9</sup> and PFGE, as these methods are broadly consistent with each other<sup>10</sup>. The authors used BEAST to infer that lineage I and II emerged in the early 19<sup>th</sup> century, while lineage III is dated to the late 19<sup>th</sup> century. Isolates from Europe are dispersed throughout the tree, and different methods of analysis provided consistent evidence for a European origin for *S. sonnei* as a whole, as well as for each of the individual lineages (Figure 1).

The stability of these four *S. sonnei* lineages points to ecological or epidemiological differences. Regression analysis of root-to-tip distance against isolation date suggests that lineage III mutates more quickly than I and II, which the authors claim might be owing to population level effects consistent with different niches. More direct evidence is provided by the non-random distribution of the non-European isolates between the different lineages. Isolates from Asia, Africa and America are largely restricted to lineage III and in particular are rarely found in lineage I. The implication is that lineage III isolates are more capable of establishing local populations subsequent to inter-continental transmission. Moreover, one particular lineage III cluster (named Global III), which emerged in the early 1970s, corresponds to almost half the isolates in the sample recovered after 1995, and is widely geographically distributed. This cluster contains even finer-scaled clusters associated with particular countries, pointing to local clonal expansion. The inter-continental transmission events out of Europe inferred by the authors are summarised in Figure 1.

### **Spread and antibiotic resistance**

Holt et al examined the distribution of known *S. sonnei* antimicrobial resistance mutations within the phylogeny, and found that global dissemination is strongly associated with multiple drug resistance (MDR). The independent acquisition of mutations in the same resistance gene within different clusters suggests strong selective pressure. Although antimicrobial treatment has little effect on *Shigella*, Holt et al speculate that the advantage conferred by resistance may operate by increasing shedding (and transmissibility) of the bacteria<sup>11</sup>, rather than simply enhanced survival in the host. The acquisition of Class 2 integrons in biotype g (lineage III) has been implicated in increased spread of *S. sonnei*, and is known to have occurred in southern Italy by the late 1980s<sup>12</sup>. The data of Holt et al, reveal three independent acquisitions of these elements within lineage III isolates during the 1960s and 1970s, immediately prior to international dissemination. It remains unclear why class 2 integrons are not found in other lineages, one possibility being the presence of potentiating mutations specific to lineage III.

The key role of antibiotic resistance genes in the global dissemination of specific *S. sonnei* clusters is underscored by limited evidence for positive selection in other genes. However, the analysis excludes the 180Kb pINV B plasmid, key in *S. sonnei* pathogenesis. This plasmid harbours the locus encoding O antigen biosynthesis, known to be under intense selection, and horizontally acquired by *S. sonnei* from *Plesiomonas shigelloides*<sup>13</sup>, a gastroenteritis causing Enterobacteria associated with poor water quality. Holt et al. argue that the cross-reactivity in O antigens between *S. sonnei* and *P. shigelloides* helps to explain the recent emergence of *S. sonnei* in transitional countries as improved sanitation would reduce

environmental immunisation induced by *P. shigelloides* infection<sup>14</sup>. Although potentially a public health conundrum, the benefits of clean drinking water will likely be of greater significance.

### Next steps

Holt et al. neatly illustrate how whole-genome sequencing of pathogens combined with phylogenetic analysis can be used to draw inferences over multiple scales, and that a small number of recently emerged clones are responsible for a disproportionately large share of the global public health burden resulting from *S. sonnei* infection. The acquisition of antibiotic resistance plays a key role in the spread of these clones, although the precise selective advantage conferred is unclear. To what degree the decline in environmental *P. shigelloides* has impacted on the spread of *S. sonnei*, and how this might possibly dovetail with the acquisition of antibiotic resistance, is also unknown. Finally, although Holt et al. raise the relevance for development of a vaccine, the implications for outbreak investigation are perhaps even more pertinent. Intercontinental travel and mass gatherings sharing a common food source<sup>15</sup> present ongoing risks for *S. sonnei* infection, and currently PFGE is the standard tool for epidemiological investigation.

- 1 Niyogi, S. K. Shigellosis. *J Microbiol* **43**, 133-143, doi:2172 [pii] (2005).
- 2 Holt, K. e. a. Shigella sonnei genome sequencing and phylogenetic analysis indicate recent global dissemination from Europe. *Nature Genetics* (2012).
- 3 Harris, S. R. *et al.* Evolution of MRSA during hospital transmission and intercontinental spread. *Science* **327**, 469-474, doi:327/5964/469 [pii] 10.1126/science.1182395 (2010).
- 4 Lan, R. & Reeves, P. R. Escherichia coli in disguise: molecular origins of Shigella. *Microbes Infect* **4**, 1125-1132, doi:S1286457902016374 [pii] (2002).
- 5 Nygren, B. L. *et al.* Foodborne outbreaks of shigellosis in the USA, 1998-2008. *Epidemiol Infect*, 1-9, doi:S0950268812000222 [pii] 10.1017/S0950268812000222 (2012).
- 6 McAdam, P. R. *et al.* Molecular tracing of the emergence, adaptation, and transmission of hospital-associated methicillin-resistant Staphylococcus aureus. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 9107-9112, doi:1202869109 [pii] 10.1073/pnas.1202869109 (2012).
- 7 Morelli, G. *et al.* Yersinia pestis genome sequencing identifies patterns of global phylogenetic diversity. *Nat Genet* **42**, 1140-1143, doi:ng.705 [pii]

10.1038/ng.705 (2010).

- 8 Shepherd, J. G., Wang, L. & Reeves, P. R. Comparison of O-antigen gene clusters of *Escherichia coli* (Shigella) *sonnei* and *Plesiomonas shigelloides* O17: *sonnei* gained its current plasmid-borne O-antigen genes from *P. shigelloides* in a recent event. *Infect Immun* **68**, 6056-6061 (2000).
- 9 Liang, S. Y. *et al.* Multilocus variable-number tandem-repeat analysis for molecular typing of *Shigella sonnei*. *Journal of clinical microbiology* **45**, 3574-3580, doi:JCM.00675-07 [pii]

10.1128/JCM.00675-07 (2007).

- 10 Chiou, C. S. *et al.* Utility of multilocus variable-number tandem-repeat analysis as a molecular tool for phylogenetic analysis of *Shigella sonnei*. *Journal of clinical microbiology* **47**, 1149-1154, doi:JCM.01607-08 [pii]

10.1128/JCM.01607-08 (2009).

- 11 Vinh, H. *et al.* Treatment of bacillary dysentery in Vietnamese children: two doses of ofloxacin versus 5-days nalidixic acid. *Trans R Soc Trop Med Hyg* **94**, 323-326 (2000).
- 12 Mammina, C., Aleo, A., Romani, C. & Nastasi, A. *Shigella sonnei* biotype G carrying class 2 integrons in southern Italy: a retrospective typing study by pulsed field gel electrophoresis. *BMC Infect Dis* **6**, 117, doi:1471-2334-6-117 [pii]

10.1186/1471-2334-6-117 (2006).

- 13 Liu, B. *et al.* Structure and genetics of *Shigella* O antigens. *FEMS Microbiol Rev* **32**, 627-653, doi:FMR114 [pii]

10.1111/j.1574-6976.2008.00114.x (2008).

- 14 Sack, D. A., Hoque, A. T., Huq, A. & Etheridge, M. Is protection against shigellosis induced by natural infection with *Plesiomonas shigelloides*? *Lancet* **343**, 1413-1415 (1994).
- 15 Lee, L. A. *et al.* An outbreak of shigellosis at an outdoor music festival. *Am J Epidemiol* **133**, 608-615 (1991).